


7D- 604608

604608


1



Approved for OTS release

6p

COPY	1	OF	1	fp
HARD COPY	\$.100			
MICROFICHE	\$.050			



The RAND Corporation
1700 MAIN ST. • SANTA MONICA • CALIFORNIA

COPYRIGHT, 1954
THE RAND CORPORATION

Analytical Approximation

Chi-Square Integral: To better than .00006 over
0 ≤ x ≤ 3 for m = 3,

$$F_m(x) = \frac{1}{2\Gamma\left(\frac{m}{2}\right)} \int_0^x \left(\frac{t}{2}\right)^{\frac{m}{2}-1} e^{-\frac{t}{2}} dt$$
$$= .26509x^{3/2} - .077320x^{5/2} + .011875x^{7/2} - .00084911x^{9/2}.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1954

Analytical Approximation

Chi-Square Integral: To better than .00001 over
 $0 \leq x \leq 2$ for $m = 2$,

$$F_m(x) = \frac{1}{2\Gamma(\frac{m}{2})} \int_0^x \left(\frac{t}{2}\right)^{\frac{m}{2}-1} e^{-\frac{t}{2}} dt$$

$$= .49977x - .12394x^2 + .019234x^3 - .0015958x^4.$$

Cecil Hastings, Jr.
 James P. Wong, Jr.
 RAND Corporation
 Copyright 1954

10-9-54

Analytical Approximation

Chi-Square Integral: To better than .0018 over
0 \leq x \leq 4 for m = 4,

$$F_m(x) = \frac{1}{2 \Gamma\left(\frac{m}{2}\right)} \int_0^x \left(\frac{t}{2}\right)^{\frac{m-1}{2}} e^{-\frac{t}{2}} dt$$
$$= .1161x^2 - .03001x^3 + .002560x^4.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1954

10-11-54

Analytical Approximation

**Chi-Square Integral: To better than .0009 over
0 ≤ x ≤ 3 for m = 3,**

$$F_m(x) = \frac{1}{2^m \left(\frac{m}{2}\right)} \int_0^x \left(\frac{t}{2}\right)^{\frac{m}{2}-1} e^{-\frac{t}{2}} dt$$

$$\approx .2589x^{3/2} - .0668x^{5/2} + .00651x^{7/2}.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1954

Analytical Approximation

Chi-Square Integral: To better than .00035 over
 $0 \leq x \leq 2$ for $m = 2$,

$$F_m(x) = \frac{1}{2 \Gamma\left(\frac{m}{2}\right)} \int_0^x \left(\frac{t}{2}\right)^{\frac{m-1}{2}} e^{-\frac{t}{2}} dt$$
$$\approx .4972x - .1163x^2 + .01207x^3.$$

Cecil Hastings, Jr.
James P. Wong, Jr.
RAND Corporation
Copyright 1954